# Figure H2. Lequesne Index

### Index of Severity for Osteoarthritis of the Knee by Lequesne et al

#### Overview:

Lequesne et al developed an index of severity for osteoarthritis for the knee (ISK). This can be used to assess the effectiveness of therapeutic interventions.

#### Sections for index:

- (1) pain or discomfort
- (2) maximum distance walked
- (3) activities of daily living

#### I Pain or Discomfort

Parameter	Finding	Points
pain or discomfort during nocturnal bedrest	none	0
	only on movement or in certain positions	1
	without movement	2
duration of morning stiffness or pain after getting up	none	0
	< 15 minutes	1
	>= 15 minutes	2
remaining standing for 30 minutes increases pain	no	0
	yes	1
pain on walking	none	0
	only after walking some distance	1
	early after starting	2
pain or discomfort after getting up from sitting without use of arms	no	0
	yes	1

#### where:

- A change in a 1991 version was to have the duration of morning stiffness scored 0 if it was 1 minute or less and 1 if it was from 1 to less than 15 minutes.
- Pain on walking in a 1991 version expanded "early after starting" to "after initial ambulation and increasingly with continued ambulation"

## II. Maximum Distance Walked

Parameter	Finding	Points
maximum distance walked	unlimited	0
	> 1 kilometer but limited	1
	about 1 kilometer (about 15 minutes)	2
	about 500 - 900 meters (about 8-15 minutes)	3
	from 300 - 500 meters	4
	from 100 - 300 meters	5
	< 100 meters	6
walking aids required	None	0
	1 walking stick or crutch	1
	2 walking sticks or crutches	2

# III. Activities of Daily Living

Parameter	Finding	Points
able to climb up a standard flight of stairs	easily	0
	with mild difficulty	0.5
	with moderate difficulty	1.0
	with marked difficulty	1.5
	impossible	2.0
able to clumb down a standard flight of stairs	easily	0
	with mild difficulty	0.5
	with moderate difficulty	1.0
	with marked difficulty	1.5
	impossible	2.0
able to squat or bend at the knee	easily	0
	with mild difficulty	0.5
	with moderate difficulty	1.0
	with marked difficulty	1.5
	impossible	2.0
able to walk on uneven ground	easily	0

with mild difficulty	0.5
with moderate difficulty	1.0
with marked difficulty	1.5
impossible	2.0

index of severity =

= SUM(points for all parameters)

Interpretation:

• minimum points for each section: 0

• maximum points for each section: 8

• minimum index score: 0

• maximum index score: 24

Index Score	Handicap	
0	none	
1 - 4	mild	
5 - 7	moderate	
8 - 10	severe	
11 - 13	very severe	
>= 14	extremely severe	

#### Modifications

The index was modified in 1997 with some minor changes to morning stiffnes and termed the "algofunctional index".

## References:

Lequesne M Mery C et al. Indexes of severity for osteoarthritis of the hip and knee. Scand J Rheumatology. 1987; Supplement 65: 85-89.

Lequesne M. Indices of severity and disease activity for osteoarthritis. Seminars in Arthritis and Rheumatism. 1991; 20 (supplement 2): 48-54.

Lequesne MG. The algofunctional indices for hip and knee osteoarthritis. J Rheumatol. 1997; 24: 779-781.